

PPl 15: The First Brown Dwarf Spectroscopic Binary

Gibor Basri

and

Eduardo L. Martín

Astronomy Department, University of California, Berkeley, CA 94720

basri@soleil.berkeley.edu, ege@popsicle.berkeley.edu

ABSTRACT

PPl 15 is the first object to have been confirmed as a brown dwarf by the lithium test (in 1995), though its inferred mass was very close to the substellar limit. It is a member of the Pleiades open cluster. Its position in a cluster color-magnitude diagram suggested that it might be binary, and preliminary indications that it is a double-lined spectroscopic binary were reported by us in 1997. Here we report on the results of a consecutive week of Keck HIRES observations of this system, which yield its orbit. It has a period of about 5.8 days, and an eccentricity of 0.4 ± 0.05 . The rotation of the stars is slow for this class of objects. Because the system luminosity is divided between 2 objects with a mass ratio of 0.85, this renders each of them an incontrovertible brown dwarf, with masses between $60\text{--}70 M_{Jup}$. We show that component B is a little redder than A by studying their wavelength-dependent line ratios, and that this variation is compatible with the mass ratio. We confirm that the system has lithium, but cannot support the original conclusion that it is depleted (which would be surprising, given the new masses). This is a system of very close objects which, if they had combined, would have produced a low mass star. We discuss the implications of this discovery for the theories of binary formation and formation of very low mass objects.

Subject headings: open clusters and associations: individual (Pleiades) — stars: low-mass, brown dwarfs — stars: binaries — stars: formation

1. Introduction

Basri, Marcy & Graham (1996; hereafter BMG) obtained in 1995 the first unambiguous detection of lithium in a brown dwarf (BD) candidate. This object was PPl 15, discovered in a CCD survey by Stauffer et al. (1994). In spite of the detection of lithium, the position of PPl 15 in the

color-magnitude diagram was very close to the expected substellar limit for the newly proposed age of the cluster (~ 115 Myr). It was possible that PPl 15 could be a very low-mass (VLM) young star or a transitional object rather than a genuine brown dwarf. BMG inferred a mass of $0.078 M_{\odot}$, which was slightly below the substellar limit at the time ($0.080 M_{\odot}$), but is slightly above the latest expected substellar limit for solar metallicity Baraffe et al. (1998, $0.075 M_{\odot}$ or $75 M_{Jup}$).

It has long been known that there is a dispersion in the Pleiades color-magnitude sequence, greater than the observational errors. For some time this was thought to be due to an age spread in the cluster, but strong arguments have been brought against that (Stauffer et al. 1995; Steele et al. 1995). It is now thought that the dispersion is likely due to unresolved binaries. These are not only expected, but explain the amplitude of the spread quite naturally. PPl 15 sits at a location which suggests that it is one of these binaries (Zapatero-Osorio et al. 1997).

An HST/WFPC2 image (J. McConnell & J. Surdej; private communication) eliminates companions to within $0.15''$ (18 AU) of the primary (with a contrast up to 5 magnitudes at I). Spectroscopically, BMG reported only that the mean line positions did not change at the level of 5 km s^{-1} (1 pixel), then coadded the spectra to analyze lithium. We re-examined BMG’s HIRES data to look for radial velocity variations using a full cross-correlation analysis of each night. To our great surprise, the cross-correlation peaks are double with comparable amplitudes, except on the first night (where there is a single peak). We thus reported that PPl 15 is a likely double-lined spectroscopic binary Basri & Martín (1998). In this paper we present a set of new HIRES observations which confirm the binarity of PPl 15 and allow us to derive its orbital parameters. We also find an $H\alpha$ flare on one of the nights, and confirm the lithium detection reported by BMG. We discuss the implications of our results for the binary frequency among BDs and the theories of binary formation.

2. Observations

Our full set of observations of PPl 15 were obtained at 3 different epochs, all at the Keck Observatory. The first 6 observations on 23-25 Nov. 1994, were the initial set in BMG which was aimed at conducting the lithium test. These were consecutive pairs of exposures on 3 consecutive nights. On 5-6 March 1995 BMG observed it four more times (again in consecutive pairs) attempting to confirm the initial preliminary detection of lithium. After analysis of these observations for binarity in early 1997 (Basri & Martín 1998), we conducted 9 observations on 1-7 Dec. 1997 to determine the orbit. In this latter run we are grateful for the cooperation of Geoff Marcy and Paul Butler, and John Stauffer and Robert O’Dell, who allowed us to trade observations on their nights for time we had allocated. It was only through this cooperation that we were able to obtain the unprecedented consecutive coverage on the Keck I telescope. The weather was not good for several nights of the run, so several spectra were obtained through clouds and barely of useable quality. Fortunately, we only need sufficient signal for a cross-correlation analysis, and can use the reddest orders. The observing log for all observations is given in Table 1.

The observations were obtained at the Keck I telescope, using the HIRES echelle spectrometer (Vogt et al. 1994). The instrumental configuration and data reduction procedure were the same as in BMG. With a dispersion of about 0.01 nm per (binned) pixel, the resolving power obtained was $R \sim 31000$. Standard echelle reductions including sky subtraction were done with our IDL routines. Fully reduced data is available about 10 minutes after the spectrum is read out, enabling decisions about whether a further exposure is needed. In addition to the spectra of PPl 15, a spectrum of Gl406 was taken with the same instrumental configuration to serve as a cross-correlation template. The star HD 93521 (O9Vp) was also observed as a relative flux standard and to measure telluric absorption line contamination.

3. Spectroscopic analysis

Each order in each PPl 15 spectrum was first median filtered to remove cosmic rays and bad columns, then a continuum fit to our flux standard was divided out and the mean spectrum normalized to unity. At this point we used our IDL routine XCORL to obtain a cross-correlation function (XCF) between that order and the corresponding one in our velocity standard. The velocity standard for Nov. 1994 and Mar. 1995 was LHS523, and for Dec. 1997 it was Gl406. A shift of up to 50 pixels in either direction produced a nice XCF with one or two sharp peaks and plenty of flat “continuum” on either side of them. Because the S/N of the spectra was generally low and the orders contain varying amounts of spectral features which can produce a correlation signal, we averaged together XCFs of the 6 redmost orders and 2 others which produced the best XCFs. We avoided orders with a strong telluric signal. This final average XCF constitutes the “observation” for each spectrum. We measured the positions of its peaks with a two-gaussian plus quadratic background fitting routine to obtain the velocity of each component in each spectrum. The errors in the individual velocity measurements are generally less than 3 km s^{-1} (though they vary with the quality of and shape of the XCF).

Because we used the same template spectrum for each observing epoch, we obtain good relative measures of the velocity shifts. The night sky airglow spectrum is also obtained in each observation, and we ensured that the template and program observation are on the same velocity scale using those. We then refer each velocity template to an absolute velocity reference by measuring the position of atomic lines in it compared with the night sky, and correcting for the solar system barycentric velocity. The systemic velocity of PPl 15 was consistent to within 2 km s^{-1} for each data set, and the overall system velocity was consistent with that published by BMG (though here we obtained a slightly higher radial velocity of $6.5 \pm 2 \text{ km s}^{-1}$ compared with their value of 4.6 km s^{-1}).

We were also able to obtain the projected rotation velocity for each star, by measuring the width of the XCF of each when they were widely separated. The method for deriving $v \sin i$ from these was the same as in Basri & Marcy (1995). The two stars appear to have the same $v \sin i$, each about 10 km s^{-1} . This is actually the lowest value (even after correction for inclination) we

have found in Pleiades VLM stars (see Oppenheimer et al. 1997), where velocities from 25-50 km s⁻¹ are more common. One possibility is that the stars have been tidally pseudo-synchronized, meaning that the angular velocity of rotation is similar to the angular velocity of the companion crossing the sky during periastron passage. The timescale for this synchronization is much shorter than for orbital circularization (which we show below has not occurred).

3.1. The masses of the components

It is important to point out that we cannot derive the actual masses from the radial velocities we have measured (as we do not know the orbital inclination). We infer the total mass of the system ($0.13M_{\odot}$) using the fact that the components are almost equal in brightness (and therefore mass), then asking what mass they would have to be based on theoretical models such as those by Baraffe et al. (1998) to match the observed luminosity and color of the system. It was a similar analysis that led BMG to the conclusion that PPl 15 was at the substellar boundary in the first place. The total system mass is now almost doubled, because each component only drops about a factor of two in luminosity and the colors remain almost the same.

There is no hope of spatially resolving the two components in the near future. Their mass ratio can be found from analysis of the individual velocities. For any arbitrary pair of phases, the mass ratio will be the ratio of the velocity difference for the primary divided by the velocity difference for the secondary. These are easily distinguished, because the 2 sets of lines are not of equal depth. We chose several pairs of phases in which the components had switched places, and measured the line positions by correlation with our velocity standards (after barycentric correction of all observations). Because of the division operation, the velocity ratio is quite sensitive to the measured centroids of the components (for example, a barycentric correction of only 1 km s⁻¹ for closely spaced observations changes that mass ratio determination by 0.15). For the phase pairs RJD 50787.9/50789.9, 49681.9/49788.8, and 49680.8/49789.9, we obtained mass ratios of 0.85, 0.91, and 0.82. We therefore set the mass ratio at 0.87 ± 0.04 from this exercise.

We can also get some information on the mass ratio from the fact that the component lines, when separated, have a ratio of line depths that is wavelength dependent. Indeed the ratio becomes more similar at longer wavelengths, as expected if the fainter component is also redder. Given our 2-gaussian fits to the XCFs in each order, we can measure the ratio of the line depths. This was done for the average of the 7 Dec 1997 spectra, and the results are given in Fig. 3. The ratio of the brighter to fainter component goes from a little over 3 at 740 nm to 1.5 at 870 nm. The line depth ratios should approximately correspond to the continuum brightness ratios. The average spectral line depths themselves should not be a function of wavelength, especially given the small expected temperature difference between the components.

To calibrate these results, we divided low dispersion spectra of Pleiades objects with spectral types M6-8 by each other. The ratios of the high resolution spectral lines between 770-790 nm




Fig. 1.— The ratio of the average line depths as a function of wavelength, as found from a cross-correlation analysis. Various echelle orders from the spectra of 7 Dec 1997 are correlated with the same orders from Gl406 (our routine XCORL produces dips rather than peaks, which look more like the average spectral lines). Both components of PP1 15 are easily visible, as is the fact that the line ratio becomes more equal at longer wavelengths.

and 810-850 nm are roughly consistent with the behavior of the divided low dispersion spectral shapes, as is the fact that the ratio is smaller for the redder wavelengths. The ratio at 750 nm is perhaps greater than expected from the low dispersion spectra. Nonetheless, the general behavior is reasonably reproduced if the brighter component has a spectral class of M6 and the fainter is M7. This is consistent with the composite spectral type of M6.5 that has been given for PPl 15 (Martín et al. 1996). The mass ratio that would be expected between those spectral types in the Pleiades is 0.8-0.9. Finally, the I magnitude difference between objects with $70M_{Jup}$ and $60M_{Jup}$ is predicted by Baraffe et al. (1998) to be 0.6 magnitudes or a factor of 1.75 in brightness, which is also about the brightness ratio found from the line depths. We therefore adopt a mass ratio of 0.85 and individual masses (based on luminosity) of $70M_{Jup}$ and $60M_{Jup}$ for PPl 15 A & B.

3.2. The orbit of PPl 15

To find the orbit we worked directly from the measured velocity differential between the two components for each night. It was obvious that the orbital period is about a week from the recurrence interval of the unsplit profiles (Basri & Martín 1998), and the fact that the velocities do not change rapidly within one night (which we tested for in Dec. 1997). We used the narrowest single profile to set the orbital epoch for $\Delta v = 0$ and then computed a set of orbits for a variety of periods in increments of 0.01 days. With initial guesses for the orbital parameters e, i, ω and an assumption of mass ratio 0.85 and total mass $0.13M_{\odot}$ (based on the system luminosity) we calculated the residual difference between the predicted and observed Δv to find at which periods it was minimized. Because of the nature of the time sampling of our data set, there are a series of minima between about 5.5-6.5 days which are not very different from each other. And of course the differences between them depend on the orbital parameters chosen. We attempted to automate a procedure for discovering the true best orbit, but found that manual exploration was able to find superior solutions. The fact is that although we can obtain very nice orbital fits with the current data set, they are apparently not sufficient for determining the unique best orbit.

The best period we could find is 5.825 days. We adopt this period, and a mass ratio of 0.85 (based on the ratio of line depths discussed above). The other orbital parameters were refined by manual searching, until the fit produced in Fig. 3.1 was found. With equal weight to all observations, the lowest residual is found for that solution (with the mean difference between the observed and predicted Δv of 2.8 km s^{-1}). Fig. 3.1 shows an alternate solution which appears as attractive (with residual of 3.1 km s^{-1}). They differ primarily in the orbital eccentricity and inclination. From tests like these, we produce the final orbital parameters listed in Table 2, where the errors reflect the uncertainty found in solutions with mean residuals that were not larger than 7 km s^{-1} . Note that the values in Table 2 do not necessarily correspond to either of the shown solutions, but are averages with errors for each parameter. We do not claim to have explored all of parameter space, nor that these orbital parameters would hold up to extensive further observations. It is unlikely, however, that they would become different enough to affect any conclusions we draw




Fig. 2.— One solution to the orbit of PP1 15. The orbital parameters are as given in Table 2, except as indicated on the Figure. This has a mean residual error in the solution for Δv of 2.8 km s^{-1} .




Fig. 3.— Another good solution to the orbit of PPl 15. It differs primarily in the eccentricity and inclination, as indicated. This has a mean residual error in the solution for Δv of 3.1 km s^{-1} .

below. The $M \sin i$ for the 2 components are $49\text{--}57M_{Jup}$ for component A and $42\text{--}48M_{Jup}$ for component B.

Our listed inclinations are based on the assumed masses from §3.1 and the orbital period – the observed velocities then imply an inclination. This is not a direct *dynamical* demonstration of the substellar nature of PPl 15. The fact that we find very reasonable orbital solutions shows that the substellar masses deduced from luminosity are consistent with the dynamical appearance of the system. What is clear is that since the system appeared to be at the substellar luminosity boundary when thought of as a single object, breaking it into two components insures that each one is definitely a brown dwarf.

3.3. The lithium abundance in PPl 15

We also reexamined the issue of lithium in PPl 15. The announcement of its detection at the AAS meeting in June 1995 (Science News 147, p.389) was the first public declaration of discovery of a brown dwarf that did not subsequently have to be recanted. BMG expressed the desire for further confirmation of the line, and of the cluster membership of PPl 15. Of course they did not realize that the object was an SB2. Proper motion confirmation of cluster membership has come from Hambly et al. (1999). Our observations of Dec. 1997 were for the purpose of finding the orbit, not detecting lithium (which is at the blue end of our spectra and requires longer exposures). Nonetheless, we can add up all the spectra, recognizing that sometimes the lithium line is single and usually is split by varying velocities, to see if there is a depression in the spectrum as expected. We find that there is, and the lithium detection is confirmed.

We can now understand better the results in BMG. On their first night the lines were not split, and the lithium feature is most obvious (they mention that 23 Nov 1994 provided the best data). In their Mar. 1995 data, not only was the moonlight contamination more severe, but the lines were quite split. In the case of very low signal-to-noise spectra, splitting the lines means essentially twice the noise for the overall lithium feature. This reduces the apparent equivalent width of the feature. The published lithium feature in BMG was also asymmetric. They thought that due merely to noise, but it turns out that modeling the line profile summed over the phases they added shows the asymmetry can actually be produced by the SB2 nature of the object. This is further confirmation that their detection was robust and that both components have lithium, but we cannot reliably measure the lithium in each component.

The accuracy of measurements is compromised by the necessity of adding spectra of various splittings (with the extra noise and continuum included). The spectrum is dominated by the hotter component, which should have an intrinsically weaker lithium line for a given abundance. Finally, the models (Allard & Hauschildt 1995) on which the depletion was based have since been revised, so the analysis should be done again. Thus, BMG’s conclusion that lithium is partially depleted in PPl 15 must be recanted; the measurements are not really good enough to say. Indeed, that

would be surprising given that the individual components are now lower in mass and should not have depleted lithium.

3.4. Variability and flaring in $H\alpha$

The $H\alpha$ line was present in each of our echelle spectra. As noted in BMG, there is a nebular contribution to the line as well as stellar emission, but this is subtracted out along with the night sky lines. Thus we have an equivalent width and line profile for each observation, with the exception of 2-4 Dec 1997 during which observing conditions were sufficiently poor that although the $H\alpha$ emission is seen, we cannot measure the underlying continuum and so cannot measure the true strength of the line. On other occasions the continuum is barely visible, so our measure of the line strength is rather inaccurate. The line varies between about 7-14 Å equivalent width on most nights; there is definitely some intrinsic variability in addition to the measurement errors (which we estimate at about 2 Å in typical cases).

On 5 Dec 1997 there is a large flare in the system. The equivalent width of the line increases by a factor of 4-5 over its typical value, and the line develops a pronounced high-velocity red wing, extending to 200 km s⁻¹. This is shown in Fig. 3.4, along with the average non-flare profiles from our 3 observing epochs. The orbital phase at this time is 0.9, so the stars are approaching each other. In addition to $H\alpha$, we see HeI (667.8 nm) and CaII (850,866 nm) emission. These lines are symmetric and quite narrow, and have equivalent widths of 2.2, 1.1, and 1.2 Å respectively. They appear to be associated with the primary only.

It is possible that the flare is associated with a magnetospheric interaction between the stars, as has been seen in RS CVn systems (UX Ari; Simon et al. 1980). Indeed, in that case red wings were also seen, extending to 450 km s⁻¹ in the MgII resonance lines. Their interpretation was in terms of a flow between the stars through interacting magnetic loops. The close T Tauri binary (DQ Tau; Basri et al. 1997) also displays a tendency to “flare” as the stars approach, but in that case it has been interpreted as more likely due to an increase in the accretion activity on those young stars. That explanation should not apply here. Of course, it is possible that the orbital phase has nothing to do with the flare; with only one event we make no claim that there is such a connection. Low mass stars are certainly known to flare even when isolated. Our observation suggests that photometric monitoring of the PPL 15 system would be a fruitful project, although we do not know if the $H\alpha$ flare was accompanied by a photometric change.




Fig. 4.— The average $H\alpha$ profiles of PPl 15 from each of the 3 epochs: Nov. 1994 (thick), Mar. 1995, Dec. 1997 (dashed, excluding the flare) are shown as the lower curves. The flare profile from 5 Dec 1997 is the much brighter line.

4. Discussion

4.1. The eccentricity of PPl 15

The tidal circularization theory of Zahn & Bouchet (1989) demonstrated that close binaries with masses ranging from 1.25 to 0.5 $M_{\odot}$ and critical periods (P_C) less than about 8 days would have circularized during their early pre-main sequence evolution. Prior to PPl 15, there were only two Pleiades binaries known with periods shorter than critical, namely HII761 ($P=3.31$ days) and HII2407 ($P=7.05$ days). Their orbits are observed to be circular (Mermilliod et al. 1992). PPl 15 has a shorter period than HII2407 yet it has a large eccentricity. According to the models of Zahn & Bouchet (1989), the value of P_C depends primarily on the pre-main sequence parameters (masses, radii and temperatures) of the binary components, because the tidal interaction is maximized for larger radii and larger convection zones (in this case the stars have remained fully convective). The fact that PPl 15 is also relatively young compared to binaries in the field is considered less relevant, since the circularization occurs essentially in the pre-main sequence phase.

For equal mass binaries with initial $e=0.3$ and components of 1.0 $M_{\odot}$ and 0.5 $M_{\odot}$, they found P_C on the main sequence of 7.80 and 7.28 days, respectively. Binaries with unequal mass components have slightly shorter critical periods. The calculations agree well with the circular orbit of HII2407 for the estimated masses of its components (0.82 $M_{\odot}$ and 0.27 $M_{\odot}$ (Mermilliod et al. 1992). Extrapolating Zahn & Bouchet’s equation 14 and using the conditions for the ‘birthline’ of deuterium burning provided by Stahler (1988), we infer that a binary with equal components of 0.1 $M_{\odot}$ has $P_C \sim 4.5$ days. The observed period of PPl 15 is longer than this value, and hence the binary is not expected to circularize. This is consistent with the eccentricity of our orbital solution.

4.2. Brown dwarf binaries

The observed frequency of very short period binaries ($P < 10$ days) is about 2% for main sequence G-dwarfs and for G- and K-type Pleiades members (Mermilliod et al. 1992). For X-ray selected stars in star forming regions the frequency of short period binaries increases by about a factor of 2 (Mathieu 1996) because of a selection bias (close binaries are brighter and more X-ray active than single stars). In flux-limited CCD surveys like that of Stauffer et al. (1994), there is also a slight bias to detect binaries because they are brighter than single stars. However, it is surprising to find that the first one selected from that type of survey has a period of only 5.8 days, because this is a rare type of binary. The distribution of binaries as a function of period has a broad peak at periods from 10^4 to 10^5 days (Duquennoy & Mayor 1991). Martín et al. (1998) resolved one binary out of six Pleiades BD candidates using HST imaging. However, we have obtained follow-up spectroscopy that does not support the membership of this binary in the Pleiades.

We have not found any binaries with separations ≥ 40 AU in a sample of 22 Pleiades BD candidates observed with HST (these observations will be presented in another paper). If the

distribution of binary frequencies among Pleiades BDs was similar to those of young stars and G dwarfs, we should have found 4.5 binaries. On the other hand, the existence of PPl 15 indicates that BD binaries may not be rare (since it was the first Pleiades BD examined for binarity). Four more pieces of evidence support this conclusion. First, Martín et al. (1999) have found one field BD binary out of two targets examined. The separation of this binary is about 5 AU. Second, the 2MASS and DENIS surveys have not reported any visual binaries among their L dwarfs (Martín et al. 1999, though there are preliminary reports of other close binaries similar to that in). Third, color-magnitude diagrams of the Pleiades VLM stars and BDs show a large spread that has been interpreted as being due mainly to unresolved binaries (Steele et al. 1993; Zapatero-Osorio et al. 1997). Fourth, the presence of an unresolved substellar secondary has been inferred from infrared spectroscopy of the VLM Pleiades star HHJ54 (Steele et al. 1995). There is no evidence for other objects in the PPl 15 system of similar mass to those in the close binary either from photometry (Zapatero-Osorio et al. 1997) or HST imaging.

If BD binaries are not rare, the fact that they have not been found at separations larger than about 40 AU in the Pleiades and the field suggests that the distribution of binary separations may be different than that of young stars and G dwarfs. One possible difference is that wide substellar binaries are less tightly bound than their stellar counterparts, and are therefore easier to disrupt by gravitational interactions. This could account for the lack of BD binaries with separations ≥ 40 AU, but it would not increase the frequency of short period binaries. The formation of PPl 15 should still be an unlikely event if the intrinsic distribution of separations is at all similar to that for stars (even if the distribution has a smaller overall scale). In order to explain the fact that PPl 15 has a very short period along with the lack of Pleiades BDs with separations ≥ 40 AU, we propose that the formation of substellar binaries tends to produce smaller separations than the formation of stellar binaries. To the best of our knowledge, this effect has not been explicitly predicted by any binary formation model.

Reviews of theories of binary formation (stellar and substellar) can be found in Bodenheimer (1998) and Bodenheimer et al. (1999). There is currently no consensus theory for the formation of very close binaries, either stellar or substellar. If they form in situ, it could depend on the inability of a collapsing core to resolve its angular momentum problem just at the last moments before it would form a single core. PPl 15 has a mass ratio close to unity, suggesting fragmentation of a bar (not too dissimilar from the now discredited fission hypotheses for binary formation). Formation of substellar objects in bars and filaments is seen in some of the calculations (Bodenheimer et al. 1999). Production of a second object in the circumstellar disk around an already formed primary is unlikely to apply to this case; the secondary is too close and too similar in mass to the primary.

Alternatively, the system we see now may not directly reflect its initial formation. Models of direct fragmentation of a collapsing cloud can produce nearly equal mass components with large eccentricities. However, the typical separations are in the range 10–100 AU (Boss 1986). If the PPl 15 system formed in a small association of protostars and was subsequently ejected from the group, the ejection velocity would have to be low because the system remains bound to the open

cluster. One possibility is that it formed as a triple substellar system, and the tight pair was produced at the expense of the ejection of the third component.

Models of multiple fragmentation yield complicated structures. Burkert & Bodenheimer (1996) have made simulations where they find an inner BD binary with a separation of about 20 AU, which induces formation of an outer BD binary with separation of about 100 AU. Such a configuration is unstable, and orbital evolution of the fragments occurs, but it has not been followed until more than 10% of the cloud mass was accreted. It is interesting to note that orbital evolution has gained great favor in explaining the orbits of the extrasolar giant planets found by radial velocity techniques (Marcy & Butler 1998).

The short period of PPl 15 is both observationally and theoretically surprising. No systematic search for other spectroscopic BD binaries has been made yet, although we and others are starting them. Explaining the formation of PPl 15 provides a new challenge for models of (substellar) binary formation. In any event, the discovery of the multiplicity of PPl 15 initiated the study of binary systems composed solely of substellar objects.

Acknowledgments: This research is based on data collected at the W. M. Keck Observatory, which is operated jointly by the University of California and California Institute of Technology. We acknowledge the support of NSF through grant AST96-18439. GB was a Miller Research Professor during this project. EM acknowledges support from the Fullbright-DGES program of the Spanish Ministry of Education. We thank again G. Marcy, P. Butler, J. Stauffer, and R. O'Dell for their cooperation during the Dec. 1997 observing run.

Table 1. Observations of PP1 15

JD(24+)	UT date	Δv	EW ($H\alpha$)	Orbital Phase
49679.92	23 Nov 1994	4.0	15	0.70
49680.81	24 Nov 1994	-33.5	7	0.85
49681.87	25 Nov 1994	-35.8	14	0.04
49788.79	12 Mar 1995	33.1	8	0.39
49789.79	13 Mar 1995	28.5	13	0.56
50783.86	01 Dec 1997	21.8	12	0.22
50784.85	02 Dec 1997	31.2	...	0.39
50785.783	03 Dec 1997	21.2	...	0.55
50785.938	03 Dec 1997	21.8	...	0.57
50786.94	04 Dec 1997	-5.0	...	0.75
50787.87	05 Dec 1997	-46.5	48	0.91
50788.83	06 Dec 1997	-17.3	15	0.07
50789.757	07 Dec 1997	29.5	11	0.23
50789.977	07 Dec 1997	31.5	11	0.27

^a Δv is the velocity separation of the two components in km s⁻¹ (a negative sign indicates the components are reversed); the emission equivalent widths of $H\alpha$ are given in Å (Dec. 2-4 were insufficiently well exposed to measure).

Table 2. Orbital Parameters for PP115

P (days)	5.825 ± 0.3
a (AU)	0.03
e	0.42 ± 0.05
i	$50^\circ \pm 5^\circ$ ^a
ω	62°
Epoch	2450782.59 ± 0.01
M_2/M_1	0.85 ± 0.05 ^b
M_1, M_2 (M_{Jup})	70, 60 (± 3) ^c

^a The inclination and semimajor axis follow from the assumed masses, which are luminosity based.

^b The mass ratio is found from spectral line depth ratios.

^cThe sum of the masses is estimated from the total system luminosity using theoretical models.

REFERENCES

- Allard, F. & Hauschildt, P.H. 1995, ApJ, 445, 433
- Basri, G., & Marcy, G. W. 1995, AJ, 109, 762
- Basri, G., Marcy, G., & Graham, J. R. 1996, ApJ, 458, 600 (BMG)
- Basri, G., Johns-Krull, C.M., Mathieu, R. 1997, AJ, 114, 781
- Basri, G., & Martín 1998, *Brown Dwarfs and Extrasolar Planets*, ASP Conf. Series Vol. 134, R. Rebolo, E. L. Martín and M. R. Zapatero Osorio (eds.), p. 284
- Baraffe, I., Chabrier, G., Allard, F., & Hauschildt, P.H. 1998, A&A, 337, 403
- Bodenheimer, P. 1998, *Brown Dwarfs and Extrasolar Planets*, ASP Conf. Series Vol. 134, R. Rebolo, E. L. Martín and M. R. Zapatero Osorio (eds.), p. 115
- Bodenheimer, P., Burkert, A., Klein, R., Boss, A. 1999, *Protostars and Planets IV*, (Mannings, Boss & Russell, eds.) (Tucson: University of Arizona Press), in press
- Boss, A. 1986, ApJS, 62, 519
- Burkert, A. & Bodenheimer, P. 1996, MNRAS, 280, 1190
- Duquennoy, A. & Mayor, M. 1991, A&A, 248, 485
- Hambly, N.C., Hodgkin, S.T., Cossburn, M.R., Jameson, R.F. 1999, MNRAS, 303, 835
- Marcy, G. & Butler, R.P. 1998, ARA&A, 36, 57
- Martín, E.L., Rebolo, R., Zapatero-Osorio, M. R. 1996, ApJ, 469, 706
- Martin, E.L., Basri, G., Brandner, W., Bouvier, J. et al. 1998, ApJ, 509, L113
- Martín, E.L., Brandner, W., Basri, G. 1999, Science, 283, 1718
- Mathieu, R. 1996, in “*The origins, evolution and destiny of binary stars in clusters*”, ASP Conf. Series Vol. 90, E. F. Milone & J. C. Mermilliod (eds.), p. 231
- Mermilliod, J.-C., Rosvick, J. M., Duquennoy, A. & Mayor, M. 1992, A&A, 265, 513
- Oppenheimer, B. R., Basri, G., Nakajima, T. & Kulkarni S. R. 1997, AJ, 113, 296
- Simon, T., Linsky, J.L., Schiffer III, F.H. 1980, ApJ, 239, 911
- Stahler, S. 1988, ApJ, 332, 804
- Stauffer, J. R., Hamilton, D. & Probst, R. G. 1994, AJ, 108, 155

- Stauffer, J. R., Liebert, J.R. & Giampapa, M. 1995, *AJ*, 109, 298
- Steele, I. A., & Jameson, R. F. 1995, *MNRAS*, 272, 630
- Steele, I. A., Jameson, R. F., & Hambly, N. C. 1993, *MNRAS*, 263, 647
- Steele, I. A., Jameson, R. F., Hodgkin, S. T. & Hambly, N. C. 1995, *MNRAS*, 275, 841
- Vogt, S.S. et al. 1994, *SPIE*, 2198, 362
- Zahn, J.-P. & Bouchet, L. 1989, *A&A*, 223, 112
- Zapatero-Osorio, M. R., Martín, E. L., & Rebolo, R. 1997, *A&A*, 323, 105